

ORIGINAL ARTICLE

Chlorine isotopic composition of perchlorate in human urine as a means of distinguishing among exposure sources

Armen Poghosyan¹, Maria Morel-Espinosa², Liza Valentin-Blasini², Benjamin C. Blount², Catterina Ferreccio³, Craig M. Steinmaus⁴ and Neil C. Sturchio^{1,5}

Perchlorate (ClO_4^-) is a ubiquitous environmental contaminant with high human exposure potential. Natural perchlorate forms in the atmosphere from where it deposits onto the surface of Earth, whereas synthetic perchlorate is manufactured as an oxidant for industrial, aerospace, and military applications. Perchlorate exposure can potentially cause adverse health effects in humans by interfering with the production of thyroid hormones through competitively blocking iodide uptake. To control and reduce perchlorate exposure, the contributions of different sources of perchlorate exposure need to be quantified. Thus, we demonstrate a novel approach for determining the contribution of different perchlorate exposure sources by quantifying stable and radioactive chlorine isotopes of perchlorate extracted from composite urine samples from two distinct populations: one in Atlanta, USA and one in Taltal, Chile (Atacama region). Urinary perchlorate from the Atlanta region resembles indigenous natural perchlorate from the western USA ($\delta^{37}\text{Cl} = +4.1 \pm 1.0\text{‰}$; $^{36}\text{Cl}/\text{Cl} = 1\,811 (\pm 136) \times 10^{-15}$), and urinary perchlorate from the Taltal, Chile region is similar to natural perchlorate in nitrate salt deposits from the Atacama Desert of northern Chile ($\delta^{37}\text{Cl} = -11.0 \pm 1.0\text{‰}$; $^{36}\text{Cl}/\text{Cl} = 254 (\pm 40) \times 10^{-15}$). Neither urinary perchlorate resembled the isotopic pattern found in synthetic perchlorate. These results indicate that natural perchlorate of regional provenance is the dominant exposure source for the two sample populations, and that chlorine isotope ratios provide a robust tool for elucidating perchlorate exposure pathways.

Journal of Exposure Science and Environmental Epidemiology advance online publication, 25 March 2015; doi:10.1038/jes.2015.18

Keywords: perchlorate; isotope; urine; chlorine

INTRODUCTION

Perchlorate (ClO_4^-) is a highly oxidized anion and is a widespread contaminant that when ingested can potentially cause adverse health effects in humans. Once in the body, perchlorate can competitively block iodide from entering the thyroid, potentially affecting further production of thyroid hormones. Deficiency of thyroid hormones has severe health effects especially for fetuses, infants, and young children. Because the thyroid hormones are crucial for neurodevelopment in the early stages of life, their deficiency can cause irreversible impairment in the nervous system, particularly in the brain.^{1–5}

Perchlorate has been detected in surface water and groundwater across the United States^{6–15} as well as in a wide range of food products including milk, infant formulas, fruits, and vegetables.^{16–22} The primary routes of perchlorate exposure to humans appear to be through oral intake of food and drinking water.^{22–26} Perchlorate has been detected in human US breast milk^{17,27–29} and in all urine samples tested as part of the National Health and Nutrition Examination Surveys between 2001 and 2008,^{30,31} indicating widespread human exposure across the United States. Urinary perchlorate is an indicator of exposure since perchlorate is not metabolized and exits the body primarily through urine.³² Children are exposed to significantly higher doses

of perchlorate compared with adults, likely because of higher rates of consumption of foods that tend to contain higher perchlorate levels.^{22,33,34}

Perchlorate originates from different natural and manufactured sources that can be distinguished by their relative isotopic signatures based on the stable chlorine isotope ratios reported in the conventional delta (δ) notation and on $^{36}\text{Cl}/\text{Cl}$ ratios. Perchlorate that forms in the atmosphere deposits onto the surface of Earth, but it accumulates mainly in dry regions such as the Atacama Desert in Chile and the Death Valley in the United States.^{35–41} Two main types of isotopically distinct natural perchlorate have been reported to date: one type having generally heavier $\delta^{37}\text{Cl}$ values (-4‰ to $+6\text{‰}$) combined with relatively high $^{36}\text{Cl}/\text{Cl}$ ratios (3100×10^{-15} to $66,500 \times 10^{-15}$) has been found at locations across the United States; and the second type characterized by anomalously low $\delta^{37}\text{Cl}$ values (-14.3‰ to -11.8‰) and relatively low $^{36}\text{Cl}/\text{Cl}$ ratios (22×10^{-15} to 590×10^{-15}) has been found only in the Atacama Desert of northern Chile.^{10,15,42,43} Atacama perchlorate is associated with nitrate-rich caliche deposits and is by far the largest known reservoir of natural perchlorate on Earth.^{8,35,44,45} Substantial quantities of Atacama perchlorate were introduced into the environment across the United States during the first half of the 20th century due to the

¹Department of Earth and Environmental Sciences, University of Illinois at Chicago, Chicago, Illinois, USA; ²Division of Laboratory Sciences, National Center for Environmental Health, Centers for Disease Control and Prevention, Atlanta, Georgia, USA; ³CENTRO FONDAP/ACCDIS, Pontificia Universidad Católica de Chile, Santiago, Chile and ⁴Arsenic Health Effects Research Program, School of Public Health, University of California, Berkeley, California, USA. Correspondence: Dr. Armen Poghosyan, Department of Earth and Environmental Sciences, University of Illinois at Chicago, 845 West Taylor Street, MC-186, Chicago, IL 60607-7059, USA.

Tel.: +1 267 207 1366.

E-mail: apogho2@uic.edu and armenpogh@hotmail.com

⁵Present address: Department of Geological Sciences, University of Delaware, Newark, Delaware 19716, USA

Received 7 November 2014; revised 20 January 2015; accepted 22 January 2015

widespread application of imported Chilean nitrate fertilizer.^{36,46–48} Synthetic perchlorate is primarily used as an oxidant in a variety of solid rocket propellants and explosives as well as in consumer products including fireworks, matches, air bags, chlorine bleach, safety flares, and chlorate herbicides.^{9,49–51} The isotopic composition of synthetic perchlorate is distinctly different from both types of natural perchlorate in its relatively consistent $\delta^{37}\text{Cl}$ value ($+0.6 \pm 1.0\%$) and its low $^{36}\text{Cl}/\text{Cl}$ ratio (1×10^{-15} to 40×10^{-15}).^{10,43,52}

Recent studies evaluated dietary sources of perchlorate in the United States by estimating overall perchlorate contribution to the food chain from all potential sources⁴⁶ or by using quantities of perchlorate in different food products, drinking water, and urine.^{22,23} However, none of these studies can differentiate between indigenous natural perchlorate, natural Atacama perchlorate, and synthetic perchlorate as the primary source of perchlorate exposure. The objective of this study was to identify the exposure source of perchlorate by measuring the chlorine isotopic composition ($\delta^{37}\text{Cl}$, $^{36}\text{Cl}/\text{Cl}$) of perchlorate extracted from urine. This novel approach can distinguish between all potential sources of perchlorate, and identify the primary source of perchlorate exposure. We hypothesized that the chlorine isotopic composition of urinary perchlorate would resemble that of ground-water, surface water, and soil in the geographical region of the population.

METHODS

Sample Collection

Perchlorate was extracted from composite urine samples for isotopic analysis by using 100 cm³ clear PVC columns filled with Purolite A530E bifunctional anion exchange resin. This resin is exceptionally selective for perchlorate,^{53–55} and similar columns have previously been used to extract perchlorate from surface water and groundwater.^{11–13,15,42,56} Detailed description of sampling column design and isotopic analytical procedures are given by Hatzinger et al.⁵⁷ The first perchlorate sample was extracted from a total volume of 40-l urine pooled from ~133 anonymously collected urines from residents of the Atlanta metro area over a 2-week collection period. The second sample was extracted from a total volume of 1.05-l urine pooled from 49 clinical residual urine samples collected from adult residents of Taltal, Chile (Atacama region). Human specimen use was in accordance with human subjects review (University of California, Berkeley). After sampling was completed, the resin columns were shipped to the University of Illinois at Chicago in a cooler filled with ice packs, and the sample columns were kept refrigerated until analysis. The amounts of perchlorate retained on each column was estimated by analyzing perchlorate concentrations in urine samples before and after the Purolite columns by using an ion chromatography tandem mass spectrometry (IC-MS/MS) at CDC.⁵⁸ Aliquots were run in triplicates for each composite sample.

Extraction of perchlorate from the resin columns was performed using procedures described by Hatzinger et al.⁵⁷ Briefly, the resin was ultrasonically cleaned in deionized water, and then it was flushed with 4-M HCl to remove common anions including organic acids. Subsequently, perchlorate was eluted with 1-M FeCl₃–4-M HCl,⁵⁵ and iron was removed by passing the diluted eluant through a cation-exchange resin, then the iron-free solution was evaporated to ~2 ml volume. The final purification of perchlorate was performed by chromatography, then perchlorate was precipitated and recrystallized as tetra-*n*-pentylammonium perchlorate (TPAClO₄).⁵⁹ The overall sample processing blank for perchlorate was 5 µg, which was < 5% of the perchlorate recovered for isotopic analysis.⁶⁰

Stable Cl Isotope Analysis ($\delta^{37}\text{Cl}$) of Perchlorate by SIMS

The stable chlorine isotope ratio ($^{37}\text{Cl}/^{35}\text{Cl}$) in perchlorate is normally analyzed by isotope ratio mass spectrometry (IRMS).^{10,11,42} However, we used secondary ion mass spectrometry (SIMS) for stable chlorine isotope ratio measurements because of the small sample size. All analyses were performed on TPAClO₄. Samples were mounted by pressing into indium metal in a 2.5-cm diameter sample holder and coating the surface with 50 nm of vapor-deposited gold. SIMS analyses were performed at the Center for Microanalysis of California Institute of Technology on an

IMS 7 f-GEO (CAMECA, Gennevilliers, France). Analyses were performed with a 10-keV Cs⁺ primary ion beam. The ion beam was focused to a diameter of ~20 µm with 5–6 nA of beam current. All isotope ratio analyses were performed by pre-sputtering of a 100 × 100 µm raster area for 120 s, and then scanning a 75 × 75 µm raster during data collection. ³⁴S contribution to ³⁵Cl results was negligible at a mass-resolving power ($M/\Delta M$) of 1200 defined at the full width at 10% of the maximum peak height; this was also confirmed under a resolution of 5000. An energy bandwidth of 45 eV was set for all SIMS measurements, and secondary ion accelerating voltage was –9 keV for all isotopic analyses. Chlorine isotopes ³⁵Cl and ³⁷Cl were measured by Faraday cups with data collection times of 1 and 2 s per cycle, respectively, for a total of 20 cycles. Data are normalized to measurements of KClO₄ isotopic reference materials USGS37 and USGS38,¹¹ which were converted to TPAClO₄ prior to measurement. Stable chlorine isotope ratios are reported in the conventional delta (δ) notation, as follows:

$$\delta^{37}\text{Cl}(\text{‰}) = \left[\left(\frac{^{37}\text{Cl}}{^{35}\text{Cl}} \right)_{\text{sample}} / \left(\frac{^{37}\text{Cl}}{^{35}\text{Cl}} \right)_{\text{SMOC}} - 1 \right] \times 1000$$

Where SMOC is the isotopic reference standard mean ocean chloride.⁶¹ Analytical precision corresponding to 1σ for $\delta^{37}\text{Cl}$ values determined from replicate measurements of reference materials is ± 1.0‰.

³⁶Cl Abundance Analysis by Accelerator Mass Spectrometry

After the analysis of the stable chlorine isotope ratios by SIMS, the remaining samples were scraped from the indium metal sample holders and diluted with USGS37 standard in TPAClO₄ form and homogenized by dissolving in 2-ml acetone. Sample J-1183 (Atlanta, USA) was diluted ~3.8 times, whereas sample J-1184 (Taltal, Chile) was diluted ~3.1 times, in order to get a larger sample size to facilitate loading of AgCl targets for accelerator mass spectrometry (AMS) measurements. Then, the diluted perchlorate samples were converted from TPAClO₄ to KClO₄ by using 0.5-mol/l KOH in absolute ethanol solution, and the KClO₄ was sealed under vacuum into 20-cm lengths of 9-mm outer diameter borosilicate glass tubing. Afterwards, KClO₄ was heated for 30 min at 600 °C in order to decompose KClO₄ according to the following reaction: $\text{KClO}_4 \rightarrow \text{KCl} + 2 \text{O}_2$ (g). Subsequently, the chloride from KCl was purified by a standard ion chromatography method developed by the Purdue Rare Isotope Measurement (PRIME) Laboratory for ³⁶Cl analyses, and then the chloride was precipitated as AgCl, which was used as a target material for the measurements of radioactive ³⁶Cl isotopic abundances by AMS. Samples for the ³⁶Cl abundances were analyzed at the PRIME Laboratory at Purdue University. The ³⁶Cl results were corrected for the instrumental background and are reported in units of $^{36}\text{Cl}/\text{Cl} \times 10^{-15}$ along with errors based on counting statistics.

Comparative Isotopic Patterns

Chlorine-36 has previously been shown to be an effective tracer of the origin of perchlorate in various environmental matrices.^{15,43} Previous work indicates unique clustering of $^{36}\text{Cl}/\text{Cl}$ ratios vs $\delta^{37}\text{Cl}$ values for perchlorate extracted from synthetic perchlorate, natural perchlorate from Atacama (Chile), and natural perchlorate from the western USA.^{42,43} The chlorine isotope pattern for synthetic perchlorate was defined based on analysis of laboratory grade perchlorate salts. The chlorine isotope pattern for natural perchlorate indigenous of the western USA was extracted from ground-water and soil samples collected from Texas, New Mexico, California, and Oregon. The chlorine isotope pattern for natural perchlorate from the Atacama Desert was characterized based on bulk soil and groundwater samples collected in that region.^{42,43} Concentration ellipses were computed using the “dataEllipse” function in the CAR (Companion to Applied Regression)⁶² package of R, an open source software environment for statistical computing and graphics,⁶³ based on a bivariate normal distribution fit to the $^{36}\text{Cl}/\text{Cl}$ ratios and $\delta^{37}\text{Cl}$ values of perchlorate reference materials. The concentration ellipses define the confidence curves within which 95% of the data is expected to lie, assuming a bivariate normality.

RESULTS

Two perchlorate samples were extracted from composite urine samples collected from geographically distinct populations located in Atlanta, USA and Taltal, Chile regions for chlorine isotopic analyses of perchlorate. A total of ~202 µg of urinary

Table 1. Stable and radioactive chlorine isotope ratios of perchlorate extracted from urine from Atlanta, USA and Taltal, Chile regions.

Sample ID	Sample location	Urine sample volume (l)	ClO_4^- on the column (μg)	Final ClO_4^- recovered (μg)	$\delta^{37}\text{Cl}_{\text{SMOC}}$ (‰)	$^{36}\text{Cl}/\text{Cl}$ (10^{-15})
J-1183	Atlanta, USA	40	198	202	$+4.1 \pm 1$	1811 ± 136
J-1184	Taltal, Chile	1.05	154	155	-11.0 ± 1	254 ± 40




Figure 1. Diagram comparing $^{36}\text{Cl}/\text{Cl}$ ratios vs $\delta^{37}\text{Cl}$ values for perchlorate extracted from urine from residents of the Atlanta, USA metro area and Taltal, Chile (Atacama region), in comparison with synthetic perchlorate, and natural perchlorate from Atacama (Chile) and the western USA.^{42,43,64} $^{36}\text{Cl}/\text{Cl}$ errors are smaller than data points. The concentration ellipses define the confidence curves within which 95% of the data is expected to lie, assuming bivariate normality.

perchlorate was recovered from the composite urine sample collected from the sample population of the Atlanta metro area, and $\sim 155 \mu\text{g}$ of urinary perchlorate was recovered from the composite urine sample collected from the residents of Taltal, Chile (Table 1). The amounts of perchlorate retained on each column and the final yields of purified perchlorate were independently determined, and the estimates agreed well for both perchlorate samples.

The urinary perchlorate samples were analyzed for the stable ($\delta^{37}\text{Cl}$) and radioactive ($^{36}\text{Cl}/\text{Cl}$) chlorine isotopic signatures in order to identify the primary exposure sources of perchlorate. The results demonstrate that the urinary perchlorate sample from the Atlanta area had a positive $\delta^{37}\text{Cl}$ value of $+4.1 \pm 1.0$ ‰ and a relatively high $^{36}\text{Cl}/\text{Cl}$ ratio of $1811 (\pm 136) \times 10^{-15}$ (Table 1). In contrast, the perchlorate sample extracted from the composite urine sample collected from multiple residents of Taltal, Chile had significantly lower stable chlorine isotopic signature ($\delta^{37}\text{Cl}$ value) of -11.0 ± 1.0 ‰ and lower radioactive chlorine isotopic abundance ($^{36}\text{Cl}/\text{Cl}$ ratio) of $254 (\pm 40) \times 10^{-15}$ compared to the urinary perchlorate from the Atlanta region.

DISCUSSION

The chlorine isotopic compositions of perchlorate samples collected from the three known principal environmental sources of perchlorate in the United States and composite urine samples from Atlanta, USA and Taltal, Chile define three distinct, non-overlapping clusters in a diagram of $^{36}\text{Cl}/\text{Cl}$ as a function of $\delta^{37}\text{Cl}$ (Figure 1). The isotopic composition of indigenous natural perchlorate from the western USA appears to be rather consistent over a wide portion of North America in terms of $\delta^{37}\text{Cl}$ values and

$^{36}\text{Cl}/\text{Cl}$ ratios. Similar isotopic signatures were observed in perchlorate from Texas, New Mexico, California, Oregon, and throughout the Great Lakes.^{15,42,43,64} Indigenous natural perchlorate samples collected from the western USA and from the Great Lakes have the highest $^{36}\text{Cl}/\text{Cl}$ ratios (3100×10^{-15} to $66,500 \times 10^{-15}$) yet reported for perchlorate, and a range of $\delta^{37}\text{Cl}$ values from -4 ‰ to $+6$ ‰. The Atacama Desert samples have much lower $^{36}\text{Cl}/\text{Cl}$ ratios (ranging from 22×10^{-15} to 590×10^{-15}) and $\delta^{37}\text{Cl}$ values (-14.3 ‰ to -11.8 ‰) compared with those of indigenous natural perchlorate from the US. Synthetic perchlorate samples have $\delta^{37}\text{Cl}$ values in the range 0.2 ‰ to 1.6 ‰ and $^{36}\text{Cl}/\text{Cl}$ ratios of 1×10^{-15} to 40×10^{-15} , which is consistent with a predominantly marine source of Cl in the brine used as for electrolytic production of perchlorate.^{43,65}

The urinary perchlorate sample from the Atlanta region more closely aligns with indigenous natural perchlorate extracted from groundwater and soil samples from the United States^{42,43} in terms of its $\delta^{37}\text{Cl}$ value and $^{36}\text{Cl}/\text{Cl}$ ratio, indicating indigenous natural perchlorate as the primary source of exposure in the sampled Atlanta residents (Figure 1). Any addition of synthetic or Atacama perchlorate would reduce the ^{36}Cl abundance in the urinary perchlorate sample from the Atlanta region, while also shifting the $\delta^{37}\text{Cl}$ result toward lower values on Figure 1 since both synthetic and Atacama perchlorates have lower $\delta^{37}\text{Cl}$ values than indigenous natural perchlorate from the United States. Overall, the shift in the $\delta^{37}\text{Cl}$ value appears to be minimal in the Atlanta sample, but the ^{36}Cl abundance is somewhat lower, compared with indigenous United States perchlorate. These features indicate negligible Atacama perchlorate contribution since it would produce a much larger shift in the $\delta^{37}\text{Cl}$ value. However, a minor synthetic perchlorate contribution to a dominantly indigenous natural source could possibly explain both the $\delta^{37}\text{Cl}$ value and the $^{36}\text{Cl}/\text{Cl}$ ratio of the Atlanta urinary perchlorate sample.

The $\delta^{37}\text{Cl}$ value and $^{36}\text{Cl}/\text{Cl}$ ratio determined for perchlorate from the urine of residents from Taltal, Chile (Atacama region) are similar to perchlorate samples collected from the Atacama region, indicating that natural Atacama perchlorate is the main source of exposure in that region (Figure 1). Perchlorate contributions from synthetic or young atmospherically produced perchlorate appear to be relatively minor, because any synthetic perchlorate contribution would increase the $\delta^{37}\text{Cl}$ value and decrease the ^{36}Cl abundance, whereas addition of significant quantities of young atmospheric perchlorate would increase the ^{36}Cl abundance while preserving the $\delta^{37}\text{Cl}$ value if we assume that the chlorine sources for atmospheric perchlorate formation in a given region have relatively constant $\delta^{37}\text{Cl}$ value over time.

Indigenous natural perchlorate samples having anomalously high $^{36}\text{Cl}/\text{Cl}$ ratios may have been affected by the presence of the bomb-produced ^{36}Cl pulse from the 1952–1962 thermonuclear tests.^{15,43} However, $^{36}\text{Cl}/\text{Cl}$ ratios $> 8000 \times 10^{-15}$ were also measured in perchlorate extracted from southwest US groundwater having ^{14}C ages of > 10 ka.⁴³ Even the lowest $^{36}\text{Cl}/\text{Cl}$ ratio of $\sim 3100 \times 10^{-15}$ in indigenous perchlorate from pre-bomb groundwater is much higher than the highest pre-bomb $^{36}\text{Cl}/\text{Cl}$ ratio of $\sim 1200 \times 10^{-15}$ in meteoric chloride near the Earth's surface across the US.^{66,67} The stratosphere is the only known location that has sufficiently high ^{36}Cl isotopic abundances⁶⁸ to account for the observed $^{36}\text{Cl}/\text{Cl}$ ratios in indigenous perchlorate samples from the western US as well as in the urinary perchlorate sample from

the Atlanta region, which strongly indicates that indigenous natural perchlorate is the dominant source of urinary perchlorate from Atlanta, USA.

The $^{36}\text{Cl}/\text{Cl}$ ratios in natural Atacama perchlorate samples are significantly lower compared to those in the indigenous natural perchlorate from the western US, which might be explained chronologically since Atacama perchlorate is much older than the western US perchlorate.^{69,70} The low $^{36}\text{Cl}/\text{Cl}$ ratios in synthetic perchlorate are consistent with chloride sources from ancient marine halite deposits.⁴³ Thus, the ^{36}Cl signature along with the stable chlorine isotope of perchlorate is a valuable tool, which can be used for tracing the source and origin of perchlorates in the environment. Although not analyzed in this study, stable oxygen isotopes ($\delta^{18}\text{O}$, $\Delta^{17}\text{O}$) of perchlorate offer additional tools for identifying perchlorate sources and origin.^{10–13,15,42,56,71}

In summary, our results demonstrate the use of chlorine isotope ratios of perchlorate extracted from pooled human urine samples for identifying the primary sources of perchlorate exposure in two geographically distinct sample populations. This novel method appears to be a robust approach, which can be used for differentiating between indigenous natural perchlorate, Atacama perchlorate and synthetic perchlorate as the primary source of perchlorate exposure, because the perchlorate molecule is stable in the environment, as well as *in vivo*,³² and can apparently retain its initial isotopic composition for centuries to millennia in groundwater and surface water systems.^{57,72} The isotopic identification of perchlorate exposure sources can aid in targeting efforts to reduce exposure and improve public health.

DISCLAIMER

The findings and conclusions in this study are those of the authors and do not necessarily represent the official views or positions of the US Centers for Disease Control and Prevention. Use of trade names and commercial sources is for identification only and does not constitute endorsement by the US Department of Health and Human Services, or the US Centers for Disease Control and Prevention.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We appreciate the assistance of Dr. Yunbin Guan and Dr. John Eiler with SIMS analyses at Caltech.

REFERENCES

- Wolff J. Perchlorate and the thyroid gland. *Pharmacol Rev* 1998; **50**: 89–105.
- Chan S, Kilby MD. Thyroid hormone and central nervous system development. *J Endocrinol* 2000; **165**: 1–8.
- National Research Council *Health implications of perchlorate ingestion*. The National Academies Press: Washington, DC, 2005.
- EPA Interim drinking water health advisory for perchlorate. *Office of Science and Technology*. US Environmental Protection Agency: Washington, DC, 2008.
- Leung AM, Pearce EN, Braverman LE. Perchlorate, iodine and the thyroid. *Best Pract Res Clin Endocrinol Metab* 2010; **24**: 133–141.
- EPA Perchlorate environmental contamination: toxicological review and risk characterization (external review draft). *National Center for Environmental Assessment*. US Environmental Protection Agency: Washington, DC, 2002.
- Urbansky ET. Perchlorate as an environmental contaminant. *Environ Sci Pollut Res Int* 2002; **9**: 187–192.
- Dasgupta PK, Martinelango PK, Jackson WA, Anderson TA, Tian K, Tock RW et al. The origin of naturally occurring perchlorate: the role of atmospheric processes. *Environ Sci Technol* 2005; **39**: 1569–1575.
- ITRC Perchlorate: overview of issues, status, and remedial options. Perchlorate-1. *Interstate Technology & Regulatory Council*. Perchlorate Team: Washington, DC, 2005.
- Sturchio NC, Bohlke JK, Gu B, Horita J, Brown GM, Beloso ADJ et al. Stable isotopic composition of chlorine and oxygen in synthetic and natural perchlorate. Gu B, Coates JD (eds) *Perchlorate: Environmental Occurrence, Interactions, and Treatment*. Springer: New York, 2006; pp 93–110.
- Bohlke JK, Hatzinger PB, Sturchio NC, Gu B, Abbene I, Mroczkowski SJ. Atacama perchlorate as an agricultural contaminant in groundwater: isotopic and chronologic evidence from Long Island, New York. *Environ Sci Technol* 2009; **43**: 5619–5625.
- Sturchio NC, Beloso A Jr., Heraty LJ, Wheatcraft S, Schumer R. Isotopic tracing of perchlorate sources in groundwater from Pomona, California. *Appl Geochem* 2014; **43**: 80–87.
- Sturchio NC, Hoaglund JR 3rd, Marroquin RJ, Beloso AD Jr., Heraty LJ, Bortz SE et al. Isotopic mapping of groundwater perchlorate plumes. *Ground Water* 2012; **50**: 94–102.
- Impellitteri CA, Saxe JP, Schmitt EC, Young KR. A survey on the temporal and spatial distribution of perchlorate in the Potomac River. *J Environ Monit* 2011; **13**: 2277–2283.
- Poghosyan A, Sturchio NC, Morrison CG, Beloso AD Jr., Guan Y, Eiler JM et al. Perchlorate in the Great Lakes: isotopic composition and origin. *Environ Sci Technol* 2014; **48**: 11146–11153.
- Jackson WA, Joseph P, Laxman P, Tan K, Smith PN, Yu L et al. Perchlorate accumulation in forage and edible vegetation. *J Agric Food Chem* 2005; **53**: 369–373.
- Kirk AB, Martinelango PK, Tian K, Dutta A, Smith EE, Dasgupta PK. Perchlorate and iodide in dairy and breast milk. *Environ Sci Technol* 2005; **39**: 2011–2017.
- Sanchez CA, Crump KS, Krieger RI, Khandaker NR, Gibbs JP. Perchlorate and nitrate in leafy vegetables of North America. *Environ Sci Technol* 2005; **39**: 9391–9397.
- Sanchez CA, Krieger RI, Khandaker N, Moore RC, Holts KC, Neidel LL. Accumulation and perchlorate exposure potential of lettuce produced in the lower Colorado River region. *J Agric Food Chem* 2005; **53**: 5479–5486.
- Sanchez CA, Krieger RI, Khandaker NR, Valentin-Blasini L, Blount BC. Potential perchlorate exposure from citrus sp. irrigated with contaminated water. *Anal Chim Acta* 2006; **567**: 33–38.
- FDA Preliminary estimation of perchlorate dietary exposure based on FDA 2004/2005 exploratory data. *Food and Drug Administration, Center for Food Safety and Applied Nutrition*. Silver Spring: Maryland, MD, 2007.
- Murray CW, Egan SK, Kim H, Beru N, Bolger PM. US Food and Drug Administration's total diet study: dietary intake of perchlorate and iodine. *J Expo Sci Environ Epidemiol* 2008; **18**: 571–580.
- Lau FK, deCastro BR, Mills-Herring L, Tao L, Valentin-Blasini L, Alwis KU et al. Urinary perchlorate as a measure of dietary and drinking water exposure in a representative sample of the United States population 2001–2008. *J Expo Sci Environ Epidemiol* 2013; **23**: 207–214.
- Mendez W, Dederick E, Cohen J. Drinking water contribution to aggregate perchlorate intake of reproductive-age women in the United States estimated by dietary intake simulation and analysis of urinary excretion data. *J Expo Sci Environ Epidemiol* 2010; **20**: 288–297.
- Huber DR, Blount BC, Mage DT, Letkiewicz FJ, Kumar A, Allen RH. Estimating perchlorate exposure from food and tap water based on US biomonitoring and occurrence data. *J Expo Sci Environ Epidemiol* 2011; **21**: 395–407.
- Blount BC, Alwis KU, Jain RB, Solomon BL, Morrow JC, Jackson WA. Perchlorate, nitrate, and iodide intake through tap water. *Environ Sci Technol* 2010; **44**: 9564–9570.
- Schier JG, Wolkin AF, Valentin-Blasini L, Belson MG, Kieszak SM, Rubin CS et al. Perchlorate exposure from infant formula and comparisons with the perchlorate reference dose. *J Expo Sci Environ Epidemiol* 2010; **20**: 281–287.
- Pearce EN, Leung AM, Blount BC, Bazrafshan HR, He X, Pino S et al. Breast milk iodine and perchlorate concentrations in lactating Boston-area women. *J Clin Endocrinol Metab* 2007; **92**: 1673–1677.
- Borjan M, Marcella S, Blount B, Greenberg M, Zhang JJ, Murphy E et al. Perchlorate exposure in lactating women in an urban community in New Jersey. *Sci Total Environ* 2011; **409**: 460–464.
- Blount BC, Pirkle JL, Osterloh JD, Valentin-Blasini L, Caldwell KL. Urinary perchlorate and thyroid hormone levels in adolescent and adult men and women living in the United States. *Environ Health Perspect* 2006; **114**: 1865–1871.
- Steinmaus C, Miller MD, Cushing L, Blount BC, Smith AH. Combined effects of perchlorate, thiocyanate, and iodine on thyroid function in the National Health and Nutrition Examination Survey 2007–08. *Environ Res* 2013; **123**: 17–24.
- Anbar M, Guttman S, Lewitus Z. The mode of action of perchlorate ions on the iodine uptake of the thyroid gland. *Int J Appl Radiat Isot* 1959; **7**: 87–96.
- Valentin-Blasini L, Blount BC, Otero-Santos S, Cao Y, Bernbaum JC, Rogan WJ. Perchlorate exposure and dose estimates in infants. *Environ Sci Technol* 2011; **45**: 4127–4132.
- Blount BC, Valentin-Blasini L, Osterloh JD, Mauldin JP, Pirkle JL. Perchlorate exposure of the US population, 2001–2002. *J Expo Sci Environ Epidemiol* 2007; **17**: 400–407.

- 35 Ericksen GE. Geology and origin of the Chilean nitrate deposits. *U.S. Geological Survey Professional Paper 1188*. United States Government Printing Office: Washington, DC, 1981.
- 36 Urbansky ET. Perchlorate chemistry: implications for analysis and remediation. *Bioremed J* 1998; **2**: 81–95.
- 37 Orris GJ, Harvey GJ, Tsui DT, Eldridge JE. Preliminary analyses for perchlorate in selected natural materials and their derivative products. *Open-file report 03-314*. U.S. Geological Survey. U.S. Dept. of the Interior: Tucson, AZ, 2003.
- 38 Plummer LN, Bohlke JK, Doughten MW. Perchlorate in Pleistocene and Holocene groundwater in north-central New Mexico. *Environ Sci Technol* 2006; **40**: 1757–1763.
- 39 Rajagopalan S, Anderson TA, Fahlquist L, Rainwater KA, Ridley M, Jackson WA. Widespread presence of naturally occurring perchlorate in high plains of Texas and New Mexico. *Environ Sci Technol* 2006; **40**: 3156–3162.
- 40 Rao B, Anderson TA, Orris GJ, Rainwater KA, Rajagopalan S, Sandvig RM et al. Widespread natural perchlorate in unsaturated zones of the southwest United States. *Environ Sci Technol* 2007; **41**: 4522–4528.
- 41 Kounaves SP, Stroble ST, Anderson RM, Moore Q, Catling DC, Douglas S et al. Discovery of natural perchlorate in the Antarctic dry valleys and its global implications. *Environ Sci Technol* 2010; **44**: 2360–2364.
- 42 Jackson WA, Bohlke JK, Gu B, Hatzinger PB, Sturchio NC. Isotopic composition and origin of indigenous natural perchlorate and co-occurring nitrate in the southwestern United States. *Environ Sci Technol* 2010; **44**: 4869–4876.
- 43 Sturchio NC, Caffee M, Beloso AD Jr., Heraty LJ, Bohlke JK, Hatzinger PB et al. Chlorine-36 as a tracer of perchlorate origin. *Environ Sci Technol* 2009; **43**: 6934–6938.
- 44 Schilt AA. *Perchloric acid and perchlorates*. G. F. Smith Chemical Co: Columbus, OH, 1979.
- 45 Ericksen GE. The Chilean nitrate deposits: the origin of the Chilean nitrate deposits, which contain a unique group of saline minerals, has provoked lively discussion for more than 100 years. *Am Scientist* 1983; **71**: 366–374.
- 46 Dasgupta PK, Dyke JV, Kirk AB, Jackson WA. Perchlorate in the United States. Analysis of relative source contributions to the food chain. *Environ Sci Technol* 2006; **40**: 6608–6614.
- 47 EPA Perchlorate environmental contamination: toxicological review and risk characterization based on emerging information (external review draft). *National Center for Environmental Assessment*. US Environmental Protection Agency: Washington, DC, 1998.
- 48 Urbansky ET, Brown SK, Magnuson ML, Kelty CA. Perchlorate levels in samples of sodium nitrate fertilizer derived from Chilean caliche. *Environ Pollut* 2001; **112**: 299–302.
- 49 Trumpolt CW, Crain M, Cullison GD, Flanagan SJP, Siegel L, Lathrop S. Perchlorate: sources, uses, and occurrences in the environment. *Remed J* 2005; **16**: 65–89.
- 50 Aziz C, Borsch R, Nicholson P, Cox E. Alternative causes of wide-spread, low concentration perchlorate impacts to groundwater. Gu B, Coates JD (eds). *Perchlorate: Environmental Occurrences, Interactions, and Treatment*. Springer: New York, 2006; pp 71–88.
- 51 Aziz CE, Hatzinger PB. Perchlorate sources, source identification, and analytical methods. Stroo HF, Ward CH (eds). *In Situ Bioremediation of Perchlorate in Groundwater*. Springer: New York, 2008; pp 55–77.
- 52 Ader M, Coleman ML, Doyle SP, Stroud M, Wakelin D. Methods for the stable isotopic analysis of chlorine in chlorate and perchlorate compounds. *Anal Chem* 2001; **73**: 4946–4950.
- 53 Batista JR, McGarvey FX, Vieira AR. The removal of perchlorate from waters using ion exchange resins. Urbansky ET (ed). *Perchlorate in the Environment*. Kluwer/Plenum: New York, 2000; pp 135–145.
- 54 Gu B, Brown GM, Chiang CC. Treatment of perchlorate-contaminated groundwater using highly selective, regenerable ion-exchange technologies. *Environ Sci Technol* 2007; **41**: 6277–6282.
- 55 Gu B, Brown GM, Maya L, Lance MJ, Moyer BA. Regeneration of perchlorate (ClO₄⁻)-loaded anion exchange resins by a novel tetrachloroferrate (FeCl₄⁻) displacement technique. *Environ Sci Technol* 2001; **35**: 3363–3368.
- 56 Bohlke JK, Sturchio NC, Gu B, Horita J, Brown GM, Jackson WA et al. Perchlorate isotope forensics. *Anal Chem* 2005; **77**: 7838–7842.
- 57 Hatzinger PB, Bohlke JK, Sturchio NC, Gu B. Guidance manual for forensic analysis of perchlorate in groundwater using chlorine and oxygen isotopic analyses. *Environmental Security Technology Certification Program (ESTCP)*. U.S. Department of Defense: Washington, DC, 2011.
- 58 Valentin-Blasini L, Blount BC, Delinsky A. Quantification of iodide and sodium-iodide symporter inhibitors in human urine using ion chromatography tandem mass spectrometry. *J Chromatogr A* 2007; **1155**: 40–46.
- 59 Dosch RG. Determination of perchlorate by precipitation with tetra-n-pentylammonium bromide. *Anal Chem* 1968; **40**: 829–831.
- 60 Poghosyan A. *Global geochemical cycle of perchlorate [Ph.D. Dissertation]*. University of Illinois at Chicago: Chicago, IL, 2013.
- 61 Godon A, Jendrzewski N, Eggenkamp HGM, Banks DA, Ader M, Coleman ML et al. A cross-calibration of chlorine isotopic measurements and suitability of seawater as the international reference material. *Chem Geol* 2004; **207**: 1–12.
- 62 Fox J, Weisberg S. *An R companion to applied regression*. 2nd SAGE Publications, Inc: Thousand Oaks, CA, 2011.
- 63 R Development Core Team R: *A language and environment for statistical computing*. R Foundation for Statistical Computing: Vienna, Austria, 2014.
- 64 Hatzinger PB, Bohlke JK, Sturchio NC, Gu B. Validation of chlorine and oxygen isotope ratio analysis to differentiate perchlorate sources and to document perchlorate biodegradation. *Final Report, ESTCP project ER-200509*. Environmental Security Technology Certification Program (ESTCP), U.S. Department of Defense: Washington, DC, 2013.
- 65 Eastoe CJ, Peryt TM, Petrychenko OY, Geisler-Cussey D. Stable chlorine isotopes in Phanerozoic evaporites. *Appl Geochem* 2007; **22**: 575–588.
- 66 Moysey S, Davis S, Zreda M, Cecil LD. The distribution of meteoric ³⁶Cl/Cl in the United States: a comparison of models. *Hydrogeol J* 2003; **11**: 615–627.
- 67 Bentley HW, Phillips FM, Davis SN. Chlorine-36 in the terrestrial environment. Fritz P, Fontes JC (eds). *Handbook of Environmental Isotope Geochemistry*. 2. Elsevier: Amsterdam, 1986; pp 427–480.
- 68 Lal D, Peters B. Cosmic ray produced radioactivity on the earth. Sitte K (ed). *Kosmische Strahlung II / Cosmic Rays II. Handbuch der Physik/Encyclopedia of Physics*. 9/46/2. Springer: Berlin, Heidelberg, 1967; pp 551–612.
- 69 Alpers CN, Brimhall GH. Middle Miocene climatic change in the Atacama desert, northern Chile: evidence from supergene mineralization at La Escondida. *Geol Soc Am Bull* 1988; **100**: 1640–1656.
- 70 Hartley AJ, Chong G. Late Pliocene age for the Atacama Desert: implications for the desertification of western South America. *Geology* 2002; **30**: 43–46.
- 71 Bao H, Gu B. Natural perchlorate has a unique oxygen isotope signature. *Environ Sci Technol* 2004; **38**: 5073–5077.
- 72 Hoering TC, Ishimori FT, McDonald HO. The oxygen exchange between oxyanions and water. II. Chlorite, chlorate and perchlorate ions. *J Am Chem Soc* 1958; **80**: 3876–3879.